



## Burn healing plants in Iranian Traditional Medicine

Sh. Fahimi<sup>1</sup>, H. Hajimehdipoor<sup>1\*</sup>, M. Abdollahi<sup>2</sup>, S.A. Mortazavi<sup>3</sup>

<sup>1</sup>*Traditional Medicine and Materia Medica Research Center and Department of Traditional Pharmacy, School of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.*

<sup>2</sup>*Department of Pharmacology and Toxicology, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran.*

<sup>3</sup>*Department of Pharmaceutics, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.*

### Abstract

Burns are known as one of the most common forms of injury with devastating consequences. Despite the discovery of several antiseptics, burn wound healing has still remained a challenge to modern medicine. Herbal products seem to possess moderate efficacy with no or less toxicity and are less expensive compared to synthetic drugs. Burn is a well-known disorder in Iranian Traditional Medicine (ITM). Iranian physicians have divided burns into various types based on the cause and recommended treatment for each type. According to ITM references, herbal therapy was the major treatment prescribed by Iranian physicians for burns. In the present study, seven ancient Iranian medical texts were screened for the herbs with burn healing effects along with their applied dosage forms. The medicinal herbs were listed and scored based on the frequency of their repetition. Moreover, the best scientific name that was suitable for each plant as well as surveying modern studies about their biological effects has been carried out. In our investigation eighteen plants with seven topical application categories have been obtained as the most frequent herbs for burn healing in ITM. Modern studies have revealed that these plants have shown some biological activities such as anti-inflammatory, antimicrobial and antioxidant effects which might establish the relationship between the mentioned activities and burn wound healing property. This list can provide a suitable resource for future researches in the field of burn treatment.

**Keywords:** anti-bacterial, anti-inflammatory, antioxidant, burn healing, Iranian Traditional Medicine

### Introduction

Burns are one of the most common and devastating forms of trauma, with outcomes spanning the spectrum from physical impairments and disabilities to emotional and mental consequences [1, 2]. Burn management entails significant duration of hospital stay, expensive medication, multiple operative

procedures and prolonged period of rehabilitation [2, 3]. Burn is defined as tissue injury caused by heat, chemicals, electricity, radiation, etc. [4]. Thermal burn and related injuries have remained a major cause of death and disability [5]. According to the depth, burn wounds are classified as first degree (superficial), second

degree (partial thickness) and third degree (full-thickness). Burn wound healing is a complicated process including inflammation due to disruption of blood vessels and extravasation of blood constituents, re-epithelialization that begins several hours after injury, formation of granulation tissue, neovascularization, migration and mitogenic stimulation of endothelial cells and finally wound contraction as a result of interaction between cells, extracellular matrix and cytokines. Severe burn wounds need to be treated as soon as possible since any delay can postpone the healing process or result in infection [4]. This makes burn care expensive and every effort should be made to provide a shorter treatment period for the burnt patients [3]. Although small burns are not usually life threatening, they need the same attention as large burns, in order to achieve functional and cosmetic outcome. Most of the early treatments include topical application of medicaments, mainly to prevent infections [5]. Healing of burn wounds still remains a challenge to modern medicine, though many antiseptics have been discovered [3]. It may be due to the fact that most of the available products have antimicrobial activity rather than wound healing effect, as well as their probable negative performance and toxicity as in the case of silver sulfadiazine on fibroblasts [4]. Improving the methods of wound healing and tissue repair offers enormous opportunities to enhance the quality of life for trauma and burn patients. It may also help to reduce health care costs. Nowadays, many drugs with natural sources have been retrieved from traditional medical systems [5]. In Iranian Traditional Medicine (ITM), plants have been used to combat various diseases and pathological conditions. Burn is a well-known disorder in ITM and several traditional prescriptions have been recommended for burn healing. According to ITM manuscripts, topical preparations have been considered as the main treatment for burns, though in some cases, phlebotomy was prescribed. ITM scholars believed that there were two stages which should be noticed in burn

treatment. The aim of the first stage was prevention of blistering by cooling and non-irritant agents. However, extra cooling should have been avoided due to the heat retention which might lead to tissue decomposition. In this situation, oils were usually useful to remove the heat. In the second stage, if the blister had been formed, detergent and desiccant medications with moderate hot and cold temperament were recommended. Generally, low fat desiccant products were preferred for wet wounds, whereas, wet temperament preparations or oils and waxes were considered as suitable medications for dry wounds [6,7]. Moreover, different dosage forms have been used in ITM for burn healing. The aim of the present study has been introducing some medicinal plants and their dosage forms used in ITM for burn healing.

## Methods

Seven ITM texts including *al-Qanun fi al-Tibb (Avicenna)* [7], *Al Havi (Rhazes)* [8], *Al-abniyah An Haghayegh el-adviah (Heravi)* [9], *al-jāmi li-mufradāt al-adwiyawa al-aghdhīy (Ibn al-Baitar)* [10], *Ekhtiarat-e-Badiee (Ansari)* [11], *Tohfāt-ul-momenin (Mo'men tonekaboni)* [12] and *Makhsan-ul-Adviah (Aghili Shirazi)* [13] were studied. The traditional word of "hargh" was considered as the keyword for extracting the plants with healing effect in all kinds of burns. First *al-Qanun fi al-Tibb*, the main text of ITM, was screened using the keyword and the burn healing property of the selected plants was checked in the rest of ITM texts to find the most frequently used plants. In addition to the temperament and used part, different dosage forms of the chosen plants, applied to burns, have been studied. So, the description of the used dosage forms has been investigated [14]. Afterwards, traditional names of the selected plants were matched to the scientific names using botanical text references [15-17]. In the next step, a substantial search has been performed in scientific databases such as "Google Scholar" and "pubmed" to find the plant biological activities concerning burn healing effect. For this

purpose, the scientific plant names in combination with the key terms including "burn healing", "wound healing", "antioxidant", "anti-inflammatory and "antibacterial" were used separately and *in vivo*, *in vitro* and clinical trials regarding to the search terms have been screened.

## Results and Discussion

Searching ITM references revealed eighteen plants as the most frequently used herbs for burn healing which have been listed in table 1. These plants have been ordered according to their repetition in the references along with their traditional names, family, temperament, used parts, dosage forms and biological effects. The plant parts which had been used for burn healing consisted of leaves, flowers, fruits, seeds, seed oil, roots, bulbs, barks and gum resins while leaves appeared to be the most used parts among others. According to table 1, most of the herbs have hot and dry temperament (>66%) while dryness is the prominent characteristic of the majority of burn healing plants introduced by Iranian scholars (>83%). Different types of dosage forms used in ITM for burn healing have been summarized in table 2. All of the mentioned dosage forms were used topically. Among them "Dohn", "Marham", "Qeirrooti" and "Zemad" (with oily base) could be suitable for dry wounds, whereas, "Tela", "Natool", "Zaroor" and "Zemad" (without oily base) were useful for wet burns. In addition to the summarized biological effects in table 1, the results of *in vitro/in vivo* tests or clinical trials performed on the plants and the probable mechanisms have been mentioned as follows:

### *Lawsonia inermis* L.

The inhibitory effects of chloroform and aqueous extracts of *L. inermis* leaves on the growth of microorganisms that are involved in burn infections has been proven [18]. The animals treated with ethanol extract of the plant leaves showed reduction in the wound area, therefore, topical formulation of ethanol extract could be successfully used for the wound healing [19,20].

The antioxidant potential of different fractions of *L. inermis* has been demonstrated through DPPH and  $\beta$ -carotene-linoleic acid assays as well [21]. The anti-inflammation effects of the plant leaves have been established in a manner that butanol and chloroform fractions exhibited more potent anti-inflammatory effect than aqueous fraction, dose dependently [22].

### *Hypericum perforatum* L.

The healing properties of *H. perforatum* ointment have been evaluated and the histopathological results, along with the wound contraction rate and period of epithelialization, have demonstrated the healing effect of *H. perforatum* ointment in linear incisions, circular excisions and thermal burn wounds [23]. Moreover, a clinical research has shown that applying a specific *H. perforatum* ointment (Gol-Daru company) three times a day for 16 days has significantly improved wound healing and has reduced scar formation compared to placebo [24]. The ethanol extract of the plant has demonstrated anti-inflammatory effects in carrageenan induced pedal edema and cotton pellet induced granuloma model [25]. A flavonoid-rich extract of *H. perforatum* L. has exhibited antioxidant activity as an effective scavenger in quenching DPPH and superoxide radicals [26]. The available literature indicates that different extracts and preparations of *H. perforatum* possess more antibacterial activity against gram-positive than gram-negative bacteria in *in vivo* and *in vitro* tests [27].

### *Myrtus communis* L.

Joursaraei *et al.* have demonstrated that *M. communis* extract could increase revascularization and the number of fibroblasts in second degree burn wounds in rats [28]. Myrtucommulone, a nonprenylated acylphloroglucinol constituent in the leaves of *M. communis*, has exerted potent *in vivo* anti-inflammatory effects [29]. In addition to the protecting effect of *M. communis* fruits against gastric ulcer, the antioxidant activity of the plant

**Table 1.** The most frequent herbs in ITM for burn healing

No.	Traditional name	phonetics	Scientific name	Family	Temperament	Part used	Dosage form	Biological effects <sup>a</sup>
1	Henna	/henna:/	<i>Lawsonia inermis</i> L.	Lythraceae	Cold and dry	Leaf	Natool	B [19, 20] , C [22], D [21], E [18]
2	Hiofariqun	/hiu:fa:ri:gu:n/	<i>Hypericum perforatum</i> L.	Hypericaceae	Hot and dry	Leaf	Zemad	A [23] B [23, 24] , C [25], D [26], E [27]
3	Ass	/a:s/	<i>Myrtus communis</i> L.	Myrtaceae	Cold and dry	Leaf/fruit	Dohn, Marham, Qeirooti, Zaroor, Zemad (with olive oil)	A [28] B [30], C [29], D [30], E [31]
4	Khobbazi	/xoobba:zi: /	<i>Malva sylvestris</i> L. <i>Malva neglecta</i> Wallr.	Malvaceae	Cold and wet	Leaf	Natool, Zemad (with olive oil)	B [32, 33], C [33, 34], D [35], E [36] B [37], C [39], D [38], E [40]
5	Zaitoon	/zæjtu:n/	<i>Olea europaea</i> L.	Oleaceae	Hot and dry (ripe fruits)	Fruit	Zemad	A [41] B [42] , C [44], D [43], E [45]
6	Semsem	/semsem/	<i>Sesamum indicum</i> L.	Pedaliaceae	Hot and wet	Seed/seed oil	Zemad	A [46] B [46], C [48] , D [47]
7	Sousan	/su:sæn/	<i>Iris sp.</i> <i>Lilium candidum</i> L.	Iridaceae Liliaceae	Hot and dry	Leaf/ bulb	Natool, Zemad (with <i>Dohn ul vard</i> ) <sup>b</sup>	C [50], D [49] C [52], D [51, 52]
8	Sanoubar	/sænoobær/	<i>Pinus pinea</i> L.	Pinaceae	Hot and dry	Bark/leaf	Tela, Zaroor	B [53], C [53], D [54], E [54]
9	Selgh	/selc/	<i>Beta vulgaris</i> L.	Amaranthaceae	Hot and dry	Leaf	Tela	C [56], D [55], E [57]
10	Ghalioon	/ga:li:u:n/	<i>Galium verum</i> L.	Rubiaceae	Hot and dry	Flower	Zemad	D [58]
11	Loban (kondor)	/looba:n/ (/koondoor/)	<i>Boswellia carteri</i> Birdw.	Burseraceae	Hot and dry	Oleogum resin	Zemad (with oily base)	C [60], D [59], E [61, 62]

**Table 1.** The most frequent herbs in ITM for burn healing (Continued)

No.	Traditional name	phonetics	Scientific name	Family	Temperament	Part used	Dosage form	Biological effects <sup>a</sup>
12	Abou khalsa	/æbu:χælsa:/	<i>Arnebia euchroma</i> (Royle) I.M.Johnst.	Boraginaceae	Hot and dry	Root	Marham, Qeirooti (with olive oil), Tela (with <i>Dohn'ul vard</i> )	A [32, 63, 64] B [65], C [64, 67], D [66], E [68]
			<i>Alkanna tinctoria</i> Taush					A [69] B [69, 70], C [73], D [71, 72], E [74]
13	Leblub	/leblab/	<i>Convolvulus arvensis</i> L.	Convolvulaceae	Dry and hot/cold <sup>c</sup>	Leaf	Natool, Qeirooti Zemad (with <i>Dohn'ul vard</i> )	D [75], E [76]
14	Narjes	/nærdʒes/	<i>Narcissus tazetta</i> L.	Amaryllidaceae	Hot and dry	Bulb	Zemad (with honey)	E [77]
15	Nil	/nil/	<i>Indigofera tinctoria</i> L.	Leguminosae	Hot and dry	Leaf	Zemad (with honey)	C [78], D [79], E [80, 81]
16	Zaimeran	/zæjmerɑ:n/	<i>Ocimum basilicum</i> L.	Lamiaceae	Hot and dry	Leaf	Tela (with rosewater), Zemad	B [82], C [84], D [83], E [85]
			<i>Ocimum minimum</i> L.					B [86], D [87], E [86]
17	Qesus	/gesu:s/	<i>Hedera helix</i> L.	Araliaceae	Hot	Flower/leaf	Qeirooti (with olive oil), Zemad	C [88, 89], D [90], E [91]
18	Quasab 'o' zarirah	/gæsæboozæri:re/	<i>Acorus calamus</i> L.	Acoraceae	Hot and dry	<sup>d</sup> -	Zemad (with <i>Dohn'ul vard</i> )	B [92], C [92, 93], D [94-96], E [93, 97]

<sup>a</sup>A: burn healing effect; B: wound healing effect; C: anti-inflammatory effect; D: antioxidant effect; E: antibacterial effect

<sup>b</sup>*Dohn'ul vard* : the oily extract of *Rosa damascena* petals

<sup>c</sup>There were different ideas about hot and cold temperament of this plant

<sup>d</sup>Not mentioned in ITM references.

has been revealed by *in vitro* studies [30]. Also, the aqueous leaf extract of *M. communis* has demonstrated an excellent antibacterial activity against *Pseudomonas aeruginosa* [31].

**Table 2.** Dosage forms used for burns in ITM

No.	Dosage form	Description
1	Dohn (/dohn/)	A dosage form prepared by extraction of the oily constituents of plant by direct compression or indirectly by using other oils especially almond, sesame or olive oil.
2	Marham (/mærhæm/)	A topical preparation consisted of an oil or wax or mixture of them especially applied for injuries and some swellings.
3	Natool (/næt u:l/)	The decoction form of plant material (with or without filtration) poured lukewarmly on the target organ where may be surrounded by a paste.
4	Qeirooti (/geiru:ti/)	A combination of oils and waxes with or without medicinal plants.
5	Tela (/tela:/)	A low viscose topical dosage form of the plant material which frequently applied on the body.
6	Zaroor (/zæru:r/)	Dried fine plant powder which sprinkled on the target organ.
7	Zemad (/zema:d/)	A soft and viscose topical preparation which may have oils and waxes and usually dressed on the location.

#### *Malva sylvestris* L.

The wound healing activity of *M. sylvestris* has been demonstrated in rat burn model by Pirbalouti *et al.* [32]. In another study, the diethyl ether extract of *M. sylvestris* flowers have effectively stimulated wound contraction and have shown increasing in well-organized bands of collagen, more fibroblasts and few inflammatory cells in the full thickness wounds of alloxan-induced diabetic rats [33]. Moreover, the ethanol extract of the plant leaves have exerted inhibitory effect on croton oil-induced ear edema in mice [34]. The antioxidant activity of *M. sylvestris* has been established by DPPH and  $\beta$ -carotene-linoleic acid assays [35]. The chloroform, ethanol and water extracts of the plant have been active against *Staphylococcus aureus* and *P. aeruginosa* [36].

#### *Malva neglecta* Wallr.

Aerial parts of *M. neglecta* have demonstrated anti-ulcerogenic activity against the ethanol-induced gastric ulcers in rat model [37]. In *in vitro* studies, hydroalcoholic extract of the plant

have shown strong antioxidant activity and reducing power by scavenging the superoxide anion, hydrogen peroxide and free radicals as well as metal chelating property compared to natural and synthetic standard antioxidants such as BHA, BHT and  $\alpha$ -tocopherol [38]. The anti-inflammatory effects of the plant extracts have been shown by Vogl *et al.* through *in vitro* studies [39]. Besides, the ethanol extract of *M. neglecta* flowers has exhibited considerable inhibitory effect against *Staphylococcus epidermidis* (MIC: 5 mg/mL) [40].

#### *Olea europaea* L.

During an *in vivo* study on full-thickness burns conducted by Edraki *et al.*, the rats treated with sea buckthorn/olive oil mixture have revealed more developed re-epithelialization and continuous basement membrane with a mature granulation tissue and have shown faster wound contraction with no infection [41]. The wound healing activity of the leaves and fruits of *O. europaea* has been studied in incision and excision models in rodents [42]. In addition, the prominent antioxidant properties of the fruit extracts of ten olive tree cultivars have been demonstrated by ABTS<sup>+</sup> method [43]. The anti-inflammatory effect of *O. europaea* has been investigated and the *n*-hexane extract of the fruits have displayed 12.7–27.8% inhibition in the carrageenan-induced hind paw edema model at 400 mg/kg [44]. Besides, the aqueous fruit extract of *O. europaea* has shown an excellent effect on *S. aureus* growth isolated from burn cases [45].

#### *Sesamum indicum* L.

The seeds and oil of *S. indicum* have shown significant reduction in period of epithelization and wound contraction in the excision and burn wound models while in the incision wound model a significant increase in the wound breaking strength has been observed [46]. Free radical scavenging capacity of antioxidants from sesame

extract has been studied in DPPH assay [47]. It has been demonstrated that sesamin, a lipid-soluble lignin isolated from *S. indicum*, has down-regulated NF- $\kappa$ B activation induced by various inflammatory stimuli. In addition, sesamin has inhibited the expression of protein COX-2 closely linked with inflammation [48].

#### *Iris sp.*

Evaluation of antioxidant effects of *I. pseudopumila* has demonstrated that in addition to the significant antiradical effect of the methanol and chloroform extracts of the flowers through DPPH assay, the chloroform extract of rhizomes and the methanol extract of the flowers have shown the most considerable activity in lipid peroxidation of liposomes [49]. Moreover, the methanol extract of *I. pseudopumila* has exhibited significant inhibitory effect on NO (an inflammatory mediator) production in macrophages [50].

#### *Lilium candidum L.*

The antioxidant capacity of *L. candidum* essential oil has been revealed by photo chemi luminescence (PCL) method [51]. The anti-inflammatory and antioxidant effects of *Pfaffia paniculata*/*Ptychopetalum olacoides* /*Lilium candidum* mixture has been measured by quantification of prostaglandin E<sub>2</sub>, leukotriene B<sub>4</sub>, histamine, and superoxide dismutase levels in an *in vitro* model of human skin culture. The results have shown a dose-dependent decrease in production of inflammatory mediators, concomitant to increased antioxidant enzyme levels under basal and lipopolysaccharide-stimulated conditions [52].

#### *Pinus pinea L.*

Süntar *et al.* have demonstrated that the essential oil obtained from the cones of *P. pinea* possessed wound healing activity in linear incision and circular excision wound models. Besides, the anti-inflammatory effect of the plant essential oil was significantly approved by the Whittle

method, which is based on the inhibition of acetic acid-induced increase in capillary permeability [53]. In addition to scavenging effect towards DPPH, the oil obtained from the nuts of *P. pinea* displayed inhibitory potentials against the isolated strains of *S. aureus* [54].

#### *Beta vulgaris L.*

The antioxidant capacity of a phenolic fraction obtained from *B. vulgaris* hydro alcoholic extract has been reported [55]. The aqueous leaves extract of the plant has shown significant anti-inflammatory activity using carrageenan induced rat paw edema method for acute inflammation and cotton pellet granuloma method for chronic inflammation [56]. Moreover, the ethanol extract obtained from the aerial parts of the plant has exhibited antibacterial effect against *S. epidermidis* and *S. aureus*. [57].

#### *Galium verum L.*

Lakić *et al.* have evaluated the antioxidant properties of methanol extract of *G. verum* in four different models and have demonstrated that the extract expressed powerful scavenging activity, reducing the DPPH and OH radical formation and neutralizing H<sub>2</sub>O<sub>2</sub>, in a dose dependent manner. The examined extract has also showed notable inhibition of lipid peroxidation [58].

#### *Boswellia carteri* Birdw.

The essential oil of *B. carteri* has shown strong antioxidant activity by inhibition of linoleic acid peroxidation [59]. Among the pentacyclic triterpenes, some boswellic acids (obtained from *Boswellia* species) are mainly responsible for many of the pharmacological effects which have been revealed to be effective in inhibition of leukotriene formation [60]. The antimicrobial activity of the essential oil isolated from the oleogum resin of *B. carteri* has been demonstrated against various microorganisms including fungi, gram-positive and gram-negative bacterial strains [61]. Moreover, binary mixture of the methanol extracts of *B. carteri* and

clarithromycine, has shown a potent synergistic effect against *P. aeruginosa* and *S. aureus* [62].

*Arnebia euchroma* (Royle) I.M.Johst.

The burn healing activity of *A. euchroma* has been revealed by Ashkani-Esfahani *et al.* They have shown that wound closure rate, fibroblast population, volume density of collagen bundles, and length density of vessels have been significantly improved by carboxymethylcellulose (CMC) gels containing *A. euchroma* hydroalcoholic extract at the concentration of 10% and 20% in third-degree burn wounds in rats [63]. In another study, it was also found that the leaves and the root extracts of *A. euchroma* had shown anti-inflammatory effect on second degree burn wounds in rats [64]. Moreover, *in vivo* healing effect of goat lipid containing *A. euchroma* root extract on burns has been demonstrated through increasing the rate of wound contraction and collagen turnover [32]. In another study, a significant wound healing activity has been observed with an ointment containing *n*-hexane root extract of *A. euchroma* (1%) in incision and excision models in mice and rats [65]. The antioxidant activity of the aqueous and methanol extracts of the plant roots has been assessed by several methods while the latter extract has shown to be more powerful in Trolox equivalent antioxidant capacity and reducing power assays [66]. The anti-inflammatory activity of the petroleum ether, chloroform, alcohol and aqueous extracts of *A. euchroma* roots has been evaluated in another research, and the results have demonstrated the prominent edema inhibition property of the extracts against carrageenan-induced rat-paw edema which were comparable to ibuprofen [67]. Besides, the ethanol extract of *A. euchroma* roots has shown interesting minimum inhibitory concentration (8 µg/mL) against *P. aeruginosa* [68].

*Alkanna tinctoria* Taush

The significant healing effect of an ointment containing alkannin and shikonin (A/S), two enantiomeric naphthoquinones from *A. tinctoria*

root extract, has been demonstrated on second intention wounds in dogs [69]. Moreover, several studies have revealed molecular and biological properties of A/S that have improve wound healing [70]. It has been declared that the dichloromethane extract of *A. tinctoria* roots, containing mainly alkannin esters, has demonstrated satisfactory antioxidant activity [71]. Besides, non-aqueous solvents and olive oil extracts of *A. tinctoria* roots with their A/S esters, have exhibited notable radical scavenging activity [72]. The efficient anti-inflammatory effect of these two natural products (A/S) has been revealed in mouse paw edema [73]. Moreover, Shikonin dyed silk exhibited bactericidal effect against *S. aureus* [74].

*Convolvulus arvensis* L.

A phenolic rich fraction of *C. arvensis* leaves, has shown antioxidant property by DPPH radical scavenging and inhibiting the oxidation of β-carotene assays [75]. In addition, the antibacterial activity of the aqueous and alcohol extracts of the plant has been demonstrated [76].

*Narcissus tazetta* L.

The plant aerial parts have shown potential antimicrobial activity against *P. aeruginosa* [77].

*Indigofera tinctoria* L.

The anti-inflammatory activity of the ethanol extract of *I. tinctoria* leaves has been evaluated by sub-planter injection of carrageenan in Wister albino rats. In this study Tyagi *et al.* has shown that oral administration of the extract, has dose dependently exhibited potent anti-inflammatory activity by reducing the induced paw edema [78]. By employing different *in vitro* assays such as reducing power, DPPH, ABTS, NO and OH radical scavenging methods, the antioxidant properties of different extracts of *I. tinctoria* leaves have been evaluated. All extracts have exhibited reducing power and strong peroxidation inhibition against linoleic acid emulsion system, although the ethyl acetate extract was found to possess more hydrogen donating ability and



higher DPPH and NO scavenging activities [79]. In a study conducted by Selvakumar and Karunakaran, *I. tinctoria* has shown higher antimicrobial activity against *P. aeruginosa*, *S. aureus* and *Bacillus subtilis* compared to penicillin and tetracycline [80]. Moreover, the chloroform extract from the plant leaves has exhibited the broadest and highest antibacterial activity compared to the butanol and ethanol extracts [81].

#### *Ocimum basilicum* L.

Using the excision wound model in rats, Salmah *et al.* have reported that the wounds treated with honey in combination with alcohol extract of *O. basilicum* have demonstrated considerable signs of dermal healing which has significantly healed earlier compared to those treated with honey alone [82]. The antioxidant property of *O. basilicum* and its effect on nitric oxide (NO) production in lipopolysaccharide-activated macrophages has been revealed [83]. Besides, *O. basilicum* methanol extract has inhibited the pro-inflammatory cytokines and mediators, which accounts for the anti-inflammatory effects [84]. Assessment of the antibacterial effects of the plant has shown that the ethanol extract of *O. basilicum* has been more effective against the pathogens such as *S. aureus* and *P. aeruginosa* when compared to the acetone extract [85].

#### *Ocimum minimum* L.

The essential oil extracted from the leaves of *O. minimum* has demonstrated anti-ulcerogenic activity in gastric ulcers induced by indomethacin and ethanol. Moreover, the essential oil has shown antibacterial activity against *S. aureus* using disc diffusion technique [86]. Lukmanul Hakkim *et al.* have revealed the antioxidant activity of the methanol leaves extract of *Ocimum* species including *O. minimum* using iron (III) reduction,  $\beta$ -carotene–linoleic acid bleaching, DPPH and superoxide anion free radical scavenging assays [87].

#### *Hedera helix* L.

In a research performed by Dumitriu *et al.*,

“Dermo HdC ”, a bioproduct obtained from the leaves of *H. helix*, has inhibited IL6 and IL8 secretion and has blocked the VEGF (Vascular Endothelial Growth Factor) which could stop the inflammation progression from acute to chronic stage [88]. Besides, the anti-inflammatory properties of the ethanol leaves extract of *H. helix* has been demonstrated using formalin-induced paw edema test in Swiss albino mice [89].  $\alpha$ -hederin and hederasaponin-C, two saponins isolated from *H. helix*, have been investigated using different antioxidant tests and have finally exhibited a strong total antioxidant activity [90]. The ethyl acetate extract of the plant has exhibited the maximum inhibitory effect against *Staphylococcus aureus*, *S. epidermidis*, *Escherichia coli*, and *Bacillus subtilis* [91].

#### *Acorus calamus* L.

Shi *et al.* have shown that in addition to the inhibitory effect on mRNA expressions of inflammatory mediators, the aqueous extracts from the fresh roots and rhizomes of *A. calamus* could significantly enhance the rate of skin wound-healing in the excisional wounding test in mice [92]. Moreover, the significant carrageenan induced anti-inflammatory property of the plant aqueous extract has been proved by several studies [93]. It has been demonstrated that the ethanol extract of *A. calamus* has exhibited hepatoprotective and antioxidant activities against acetaminophen-induced liver injury in rats which are comparable to the standard drug, silymarin [94]. In addition to the of ethanol extract of *A. calamus* rhizomes [95], the leaf methanol extract has exhibited remarkable DPPH radical-scavenging activity, ability of chelating ferrous ions and reducing power, whereas the rhizome extract has displayed the strongest superoxide anion-scavenging activity as well as radical scavenging ability [95,96]. Besides, the antioxidant activity of *A. calamus* leaves and rhizomes extracts were comparable to the standard antioxidants [96]. The alcohol extract and the essential oil of *A. calamus* have represented potent *in vitro* activity against some pathogenic and nonpathogenic fungi, Gram

negative and Gram positive bacteria specially against *S. aureus* [93]. In addition, the antibacterial effect of the plant leaf extract against *P. aeruginosa* has been demonstrated [97].

### Conclusion

Wound healing is a complicated process occurring in injured tissues to restore its construction and return the damaged tissue to its normal situation as soon as possible [98]. The aim of the healing process is prevention of pathogens invasion, confirm the integrity of damaged tissue, and reconstruct the skin physiological function [99]. There are different factors involved in burn healing process; oxidants are considered as inhibitory factors to wound healing so, protecting the tissue from oxidative damage is the important role of antioxidants which leads to survival of ischemic skin flaps or improvement of wound healing [100]. On the other hand, the excessive and unbalanced inflammation could delay healing and enhance scarring which suggests a promising target for future therapeutic interventions or even predisposed tissue to cancer development [101-103]. Therefore, the anti-inflammatory compounds could act as effective agents in wound healing [104]. Despite the natural progression of the burn healing process, infections can delay the process by several mechanisms. Hence, infection is the major complication of burn injuries and antibacterial agents play a critical role in the burn healing process [102,104-107]. Plants have an extensive potential for management and treatment of wounds and burns with their antioxidant, anti-inflammatory and antimicrobial activities due to their phytochemical constituents such as alkaloids, essential oils, flavonoids, tannins, terpenoids, saponins, fatty acids and phenolic compounds [4,106,108]. Low cost, availability and fewer side effects are other advantages of herbal remedies [4]. In the present investigation, we have introduced eighteen plants which were the most frequent herbs in ITM for burn healing.

Modern studies have also shown that all of the mentioned plants possessed at least one biological effect corresponding to burn healing activity while antioxidant property was the prominent characteristic of the burn healing herbs. Moreover, among the plants which have been evaluated for wound healing effect, only five plants including “Hiofariqun”, “Ass”, “Zaitoon”, “Semsem” and “Abou khalsa” demonstrated burn healing property. Hence, the rest of the introduced herbs could be appropriate candidates for future burn healing studies and alternative medicaments for management of burn wounds.

### Acknowledgments

The article was based on Ph.D. thesis of Traditional Pharmacy (Shirin Fahimi, No. 128), granted by School of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran (No. 111).

### Declaration of interest

The authors declare that there is no conflict of interest. The authors alone are responsible for the content of the paper.

### References

- [1] Ezzati A, Bayat M, Taheri S, Mohsenifar Z. Low-level laser therapy with pulsed infrared laser accelerates third -degree burn healing process in rats. *J Reh Res Dev*. 2009; 46(4): 543-554.
- [2] Upadhyay NK, Kumar R, Siddiqui MS, Gupta A. Mechanism of wound-healing activity of *Hippophae rhamnoides* L. leaf extract in experimental burns. *Evid-Based Complemen Altern Med*. 2011; 1-9.
- [3] Khorasani G, Hosseinimehr SJ, Zamani P, Ghasemi M, Ahmadi A. The effect of Saffron (*Crocus Sativus*) extract for healing of second-degree burn wound in Rats. *Keio J Med*. 2008; 57(4): 190-195.
- [4] Bahramsoltani R, Farzaei MH, Rahimi R. Medicinal plants and their natural components as future drugs for the treatment

- of burn wounds: an integrative review. *Arch Dermatol Res.* 2014; 306(7): 601-617.
- [5] Srivastava P, Durgaprasad S. Burn wound healing property of *Cocos nucifera*: An appraisal. *Indian J Pharmacol.* 2008; 40(4): 144-146.
- [6] Nasiri E, Hosseinimehr SJ, Azadbakht M, Madani SA. A review of natural products for burn healing based on the Iranian Traditional Medicine. *J Mazand Univ Med Sci* 2014; 23(110): 263-276.
- [7] Avicenna. *Al-qanun fi al -tibb.* Beirut: Dar Ehia Al-Tourath Al-Arabi, 2005 (in Arabic).
- [8] Rhazes. *Al havi.* 1<sup>st</sup> ed. Afsharypuor S, (Trans.) Tehran: Academy of Medical Sciences, 2005 (in Persian).
- [9] Heravi M. *Al-abniah an haghayegh el-adviah* 1<sup>st</sup> ed. Tehran: Tehran University, 1967 (in Persian).
- [10] Ibn al-Baitar A. *Al-jāmi li-mufradāt al-adwiyawa al-aghdiyya.* 1<sup>st</sup> ed. Tehran: Iran University of Medical Sciences, Research Institute for Islamic and Complementary Medicine, 2008 (in Arabic).
- [11] Ansari A. *Ekhtiarat-e-badiee.* 1<sup>st</sup> ed. Tehran: Razi Pharmaceutical Distribution Company Press, 1992 (in Persian).
- [12] Tonekaboni M. *Tohfat-ul-momenin.* 1<sup>st</sup> ed. Tehran: Traditional Medicine and Materia Medica Research Center of Shahid Beheshti University of Medical Sciences and Nashr-e-shahr Press, 2007 (in Persian).
- [13] Aghili Shirazi MH. *Makhzan-ul-adviah.* 1<sup>st</sup> ed. Tehran: Iran University of Medical Sciences, Research Institute for Islamic and Complementary Medicine, 2008 (in Persian).
- [14] Afsharypuor S, Shams Ardekani M, Mosaddegh M, Ghannadi A, Mohagheghzadeh A, Badr P. *Introduction to Iranian traditional pharmacy and pharmaceutical dosage forms.* 1<sup>st</sup> ed. Tehran: Choogan press and Faculty of Traditional Medicine, Tehran University of Medical sciences, 2013 (in Persian).
- [15] Soltani A. *Encyclopedia of traditional medicine, medicinal plants.* 1<sup>st</sup> ed. Tehran: Iran University of Medical Sciences and Health Services, Institute of Medical History, Islamic and Complementary Medicine Research, Medicinal Plants Group, 2004 (in Persian).
- [16] Amin Gh. *The most common medicinal plants of Iran.* 1<sup>st</sup> ed. Tehran: Tehran University of Medical Sciences and Health Services, 2005 (In Persian).
- [17] Ghahraman A, Okhovvat AR. *Matching the old medicinal plant names with scientific terminology.* Vol 1. 1<sup>st</sup> ed. Tehran: Tehran University, 2004 (in Persian).
- [18] Muhammad HS, Muhammad S. The use of *Lawsonia inermis* Linn. (henna) in the management of burn wound infections. *Afr J biotechnol.* 2005; 4(9): 934-937.
- [19] Agarwal P, Alok S, Verma A. An update on Ayurvedic herb Henna (*Lawsonia inermis* L.): a review. *Int J Pharm Sci Res.* 2014; 5(2): 330-339.
- [20] Alam G, Singh MP, Singh A. Wound healing potential of some medicinal plants. *Int J Pharm Sci Rev Res.* 2011; 9(1): 136-145.
- [21] Hsouna AB, Trigui M, Culioli G, Blache Y, Jaoua S. Antioxidant constituents from *Lawsonia inermis* leaves: isolation, structure elucidation and antioxidative capacity. *Food Chem.* 2011; 125(1): 193-200.
- [22] Alia B, Bashir A, Tanira M. Anti-inflammatory, antipyretic, and analgesic effects of *Lawsonia inermis* L. (henna) in rats. *Pharmacology.* 1995; 51(6): 356-363.
- [23] Prisacaru AI, Andritoiu CV, Andriescu C, Havarneanu EC, Popa M, Motoc AG, Sava A. Evaluation of the wound-healing effect of a novel *Hypericum perforatum* ointment in skin injury. *Rom J Morphol Embryol.* 2013; 54(4): 1053-1059.
- [24] Vujanovic S, Vujanovic J. Bioresources in the pharmacotherapy and healing of burns: a mini-review. *Burns.* 2013; 39(6): 1031-1038.
- [25] Kumar V, Singh PN, Bhattacharya SK. Anti-inflammatory and analgesic activity of Indian *Hypericum perforatum* L. *Indian J*

- Exp Biol.* 2001; 39(4): 339-343.
- [26] Zou Y, Lu Y, Wei D. Antioxidant activity of a flavonoid-rich extract of *Hypericum perforatum* L. *in vitro*. *J Agric Food Chem.* 2004; 52(16): 5032-5039.
- [27] Saddiqe Z, Naeem I, Maimoona A. A review of the antibacterial activity of *Hypericum perforatum* L. *J Ethnopharmacol.* 2010; 131(3): 511-521.
- [28] Joursaraei S, Moghadamnia A, Rahmani firouzjaei A, Miri S, Omranirad A, Saghebi R, Hashemi S. A comparison on histopathological effects of Myrtle extract and silver sulfadiazine 1% on healing of second degree burn wound in rats. *The J Qazvin Univ Med Sci.* 2006; 10(1): 5-15.
- [29] Rossi A, Di Paola R, Mazzon E, Genovese T, Caminiti R, Bramanti P, Pergola C, Koeberle A, Werz O, Sautebin L. Myrtucommulone from *Myrtus communis* exhibits potent anti-inflammatory effectiveness *in vivo*. *J Pharm Exp Ther.* 2009; 329(1): 76-86.
- [30] Farzaei MH, Rahimi R, Abbasabadi Z, Abdollahi M. An evidence-based review on medicinal plants used for the treatment of peptic ulcer in traditional Iranian medicine. *Int J Pharmacol.* 2013; 9(2): 108-124.
- [31] Al-Saimary IE, Bakr SS, Jaffar T, Al-Saimary AE, Salim H, Al-Muosawi R. Effects of some plant extracts and antibiotics on *Pseudomonas aeruginosa* isolated from various burn cases. *Saudi Med J.* 2002; 23(7): 802-805.
- [32] Pirbalouti AG, Yousefi M, Nazari H, Karimi I, Koohpayeh A. Evaluation of burn healing properties of *Arnebia euchroma* and *Malva sylvestris*. *Electron J Biol.* 2009; 5(3): 62-66.
- [33] Pirbalouti AG, Azizi S, Koohpayeh A, Hamedi B. Wound healing activity of *Malva sylvestris* and *Punica granatum* in alloxan-induced diabetic rats. *Acta Pol Pharm.* 2010; 67(5): 511-516.
- [34] Confortia F, Sosa S, Marrelli M, Menichini F, Statti GA, Uzunov D, Tubaro A, Menichini F, Loggia RD. *In vivo* anti-inflammatory and *in vitro* antioxidant activities of Mediterranean dietary plants. *J Ethnopharmacol.* 2008; 116: 144-151.
- [35] Gasparetto JC, Martins CA, Hayash SS, Otuky MF, Pontarolo R. Ethnobotanical and scientific aspects of *Malva sylvestris* L.: a millennial herbal medicine. *J Pharm Pharmacol.* 2012; 64(2): 172-189.
- [36] Bonjar S. Evaluation of antibacterial properties of some medicinal plants used in Iran. *J Ethnopharmacol.* 2004; 94(2): 301-305.
- [37] Gürbüz I, Özkan AM, Yesilada E, Kutsal O. Anti-ulcerogenic activity of some plants used in folk medicine of Pinarbasi (Kayseri, Turkey). *J Ethnopharmacol.* 2005; 101(1): 313-318.
- [38] Güder A, Korkmaz H. Evaluation of *in vitro* antioxidant properties of hydroalcoholic solution extracts *Urtica dioica* L., *Malva neglecta* Wallr. and their mixture. *Iran J Pharm Res.* 2012; 11(3): 913-923.
- [39] Vogl S, Picker P, Mihaly-Bison J, Fakhrudin N, Atanasov AG, Heiss EH, Wawrosch C, Reznicek G, Dirsch VM, Saukel J. Ethnopharmacological *in vitro* studies on Austria's folk medicine—An unexplored lore *in vitro* anti-inflammatory activities of 71 Austrian traditional herbal drugs. *J Ethnopharmacol.* 2013; 149(3): 750-771.
- [40] Seyyednejad SM, Koochak H, Darabpour E, Motamedi H. A survey on *Hibiscus rosa-sinensis*, *Alcea rosea* L. and *Malva neglecta* Wallr as antibacterial agents. *Asian Pac J Trop Med.* 2010; 3(5): 351-355.
- [41] Edraki M, Akbarzadeh A, Hosseinzadeh M, Tanideh N, Salehi A, Koohi-Hosseinabadi O. Healing effect of sea buckthorn, olive oil and their mixture on full-thickness burn wounds. *Adv Skin Wound Care.* 2014; 27(7): 317-323.
- [42] Ghosh PK, Gaba A. Phyto-extracts in wound healing. *Int J Pharm Pharm Sci.* 2013; 16(5): 760-820.
- [43] Silva S, Gomes L, Leitao F, Coelho A, Boas

- LV. Phenolic compounds and antioxidant activity of *Olea europaea* L. fruits and leaves. *Food Sci Technol Int.* 2006; 12(5): 385-395.
- [44] Süntar İP, Akkol EK, Baykal T. Assessment of anti-inflammatory and antinociceptive activities of *Olea europaea* L. *J Med Food.* 2010; 13(2): 352-356.
- [45] Alsaimary IE. Efficacy of some antibacterial agents on *Staphylococcus aureus* isolated from various burn cases. *Int J Med Sci.* 2009; 1(4): 110-114.
- [46] Kiran K, Asad M. Wound healing activity of *Sesamum indicum* L. seed and oil in rats. *Indian J Exp Biol.* 2008; 46(11): 777-782.
- [47] Suja KP, Jayalekshmy A, Arumughan C. Free radical scavenging behavior of antioxidant compounds of sesame (*Sesamum indicum* L.) in DPPH system. *J Agric Food Chem.* 2004; 52(4): 912-915.
- [48] Harikumar KB, Sung B, Tharakan ST, Pandey MK, Joy B, Guha S, Krishnan S, Aggarwal BB. Sesamin manifests chemopreventive effects through the suppression of NF- $\kappa$ B-regulated cell survival, proliferation, invasion, and angiogenic gene products. *Mol Cancer Res.* 2010; 8(5): 751-761.
- [49] Rigano D, Conforti F, Formisano C, Menichini F, Senatore F. Comparative free radical scavenging potential and cytotoxicity of different extracts from *Iris pseudopumila* Tineo flowers and rhizomes. *Nat prod res.* 2009; 23(1): 17-25.
- [50] Conforti F, Rigano D, Menichini F, Loizzo MR, Senatore F. Protection against neurodegenerative diseases of *Iris pseudopumila* extracts and their constituents. *Fitoterapia.* 2009; 80(1): 62-67.
- [51] Bulut C, Altıok E, Bayraktar O, Ülkü S. Antioxidative and antimicrobial screening of 19 commercial essential oils in Turkey. 1<sup>st</sup> International Medicinal and Aromatic Plants Conference on Culinary Herbs. 2007 April 29 - May 4; Antalya, Turkey.
- [52] Eberlin S, Del Carmen Velazquez Pereda M, de Campos Dieamant G, Nogueira C, Werka RM, De Souza Queiroz ML. Effects of a Brazilian herbal compound as a cosmetic eyecare for periorbital hyperchromia ("dark circles"). *J Cosmet Dermatol.* 2009; 8(2): 127-135.
- [53] Süntar I, Tumen I, Ustün O, Keleş H, Küpeli Akkol E. Appraisal on the wound healing and anti-inflammatory activities of the essential oils obtained from the cones and needles of *Pinus* species by *in vivo* and *in vitro* experimental models. *J Ethnopharmacol.* 2012; 139(2): 533-540.
- [54] Orhan I, Özçelik B, Şener B. Evaluation of antibacterial, antifungal, antiviral, and antioxidant potentials of some edible oils and their fatty acid profiles. *Turk J Biol.* 2011; 35(2): 251-258.
- [55] Ninfali P, Bacchiocca M, Antonelli A, Biagiotti E, Di Gioacchino A, Piccoli G, Stocchi V, Brandi G. Characterization and biological activity of the main flavonoids from Swiss Chard (*Beta vulgaris* subspecies *cycla*). *Phytomedicine.* 2007; 14(2): 216-221.
- [56] Jain S, Garg VK, Sharma PK. Anti-inflammatory activity of aqueous extract of *Beta vulgaris* L. *J Basic Clin Pharm.* 2011; 2(2): 83-86.
- [57] Koochak H, Seyyednejad SM, Motamedi H. Preliminary study on the antibacterial activity of some medicinal plants of Khuzestan (Iran). *Asian Pac J Trop Med.* 2010; 3(3): 180-184.
- [58] Lakić NS, Mimica-Dukić NM, Isak JM, Božin BN. Antioxidant properties of *Galium verum* L. (Rubiaceae) extracts. *Cent Eur J Biol.* 2010; 5(3): 331-337.
- [59] Prakash B, Mishra PK, Kedia A, Dubey N. Antifungal, antiaflatoxin and antioxidant potential of chemically characterized *Boswellia carterii* Birdw essential oil and its *in vivo* practical applicability in preservation of *Piper nigrum* L. fruits. *LWT-Food Sci Technol.* 2014; 56(2): 240-247.
- [60] Ammon H. Boswellic acids in chronic inflammatory diseases. *Planta Med.* 2006;

- 72(12): 1100-1116.
- [61] Rahimi R, Shams-Ardekani MR, Abdollahi M. A review of the efficacy of traditional Iranian medicine for inflammatory bowel disease. *World J Gastroenterol.* 2010; 16(36): 4504-4514.
- [62] Al-kuraishy HM, Al-gareeb AI, Ashoor AW, Al-windy SA. Modifying effects of *Boswellia carteri* on clarithromycine action: *In vitro* antibacterial study against common sensitive bacterial strains. *Int J Drug Dev Res.* 2012; 4(3): 155-162.
- [63] Ashkani-Esfahani S, Imanieh MH, Meshksar A, Khoshneviszadeh M, Noorafshan A, Geramizadeh B, Ebrahimi S, Handjani F, Nadimi E, Jafari SMS. Enhancement of fibroblast proliferation, vascularization and collagen synthesis in the healing process of third-degree burn wounds by topical *Arnebia euchroma*, a herbal medicine. *Galen Med J.* 2013; 1(2): 53-59.
- [64] Ashkani-Esfahani S, Imanieh M, Khoshneviszadeh M, Meshksar A, Noorafshan A, Geramizadeh B, Ebrahimi S, Handjani F, Tanideh N. The healing effect of *Arnebia euchroma* in second degree burn wounds in rat as an animal model. *Iran Red Crescent Med J.* 2012; 14(2): 70-74.
- [65] Akkol EK, Koca U, Peşin I, Yılmaz D, Tokar G, Yeşilada E. Exploring the wound healing activity of *Arnebia densiflora* (Nordm.) Ledeb. by *in vivo* models. *J Ethnopharmacol.* 2009; 124(1): 137-141.
- [66] Chang MJ, Huang GJ, Ho YL, Lin IH, Huang SS, Chang TN, Chang HY, Chang YS. Study on the antioxidant activities of crude extracts from the roots of *Arnebia euchroma* and *Lithospermum erythrorhizon*. *Mid Taiwan J Med.* 2008; 13: 113-121.
- [67] Kaith B, Kaith N, Chauhan N. Anti-inflammatory effect of *Arnebia euchroma* root extracts in rats. *J Ethnopharmacol.* 1996; 55(1): 77-80.
- [68] Kumar GP, Singh SB. Antibacterial and antioxidant activities of ethanol extracts from trans Himalayan medicinal plants. *Eur J Appl Sci.* 2011; 3(2): 53-57.
- [69] Karayannopoulou M, Tsioli V, Loukopoulos P, Anagnostou TL, Giannakas N, Savvas I, Papazoglou LG, Kaldrymidou E. Evaluation of the effectiveness of an ointment based on alkannins/shikonins on second intention wound healing in the dog. *Can J Vet Res.* 2011; 75(1): 42-48.
- [70] Papageorgiou V, Assimopoulou A, Ballis A. Alkannins and shikonins: a new class of wound healing agents. *Curr Med Chem.* 2008; 15(30): 3248-3267.
- [71] Assimopoulou A, Boskou D, Papageorgiou V. Antioxidant activities of alkannin, shikonin and *Alkanna tinctoria* root extracts in oil substrates. *Food Chem.* 2004; 87(3): 433-438.
- [72] Assimopoulou A, Papageorgiou V. Radical scavenging activity of *Alkanna tinctoria* root extracts and their main constituents, hydroxynaphthoquinones. *Phytother Res.* 2005; 19(2): 141-147.
- [73] Kourounakis AP, Assimopoulou AN, Papageorgiou VP, Gavalas A, Kourounakis PN. Alkannin and Shikonin: effect on free radical processes and on inflammation-a preliminary pharmacochemical investigation. *Archiv Der Pharmazie.* 2002; 335(6): 262-266.
- [74] Dhandapani R, Sarkar AK. Antibacterial activity and UV property of shikonin on silk substrate. *JTATM.* 2007; 5(4): 1-7.
- [75] Elzaawely A, Tawata S. Antioxidant activity of phenolic rich fraction obtained from *Convolvulus arvensis* L. leaves grown in Egypt. *Asian J Crop Sci.* 2012; 4(1): 32-40.
- [76] Kaur M, Kalia A. *Convolvulus arvensis*: A useful weed. *Int J Pharm Pharm Sci.* 2012; 4(1): 38-40.
- [77] Talib WH, Mahasneh AM. Antimicrobial, cytotoxicity and phytochemical screening of Jordanian plants used in traditional medicine. *Molecules.* 2010; 15(3): 1811-1824.
- [78] Tyagi PK, Rai VK, Pahria AK, Kumar SS, Singh Y, Sharma M, Goyal M. Preliminary

- phytochemical screening and evaluation of anti-inflammatory activity of ethanolic extract of leaves of *Indigofera tinctoria* Linn. *J Curr Pharm Res.* 2010; 3: 47-50.
- [79] Anusuya N, Manian S. Antioxidant and free radical scavenging potential of different solvent extracts of *Indigofera tinctoria* L. leaves. *Int J Pharm Pharm Sci.* 2013; 5(1): 142-147.
- [80] Selvakumar S, Karunakaran C. Antimicrobial efficacy of *Senna auriculata*, *Pongamia glabra* and *Indigofera tinctoria* against pathogenic Microorganisms. *Int J Pharm Tech Res.* 2010; 2(3): 2054-2059.
- [81] Ravichandran R, Alwarsamy M. Phytochemical screening and *in-vitro* antibacterial activity of leaf extracts of *Indigofera tinctoria* Linn. *Int J Adv Pharm Res.* 2012; 3(4): 872-877.
- [82] Salmah I, Mahmood A, Sidik K. Synergistic effects of alcoholic extract of sweet basil (*Ocimum basilicum* L.) leaves and honey on cutaneous wound healing in rats. *Int J Mol Med Adv Sci.* 2005; 1(3): 220-224.
- [83] Abas F, Lajis NH, Israf D, Khozirah S, Umi Kalsom Y. Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chem.* 2006; 95(4): 566-573.
- [84] Selvakkumar C, Gayathri B, Vinaykumar KS, Lakshmi BS, Balakrishnan A. Potential anti-inflammatory properties of crude alcoholic extract of *Ocimum basilicum* L. in human peripheral blood mononuclear cells. *J Health Sci.* 2007; 53(4): 500-505.
- [85] Durga KR, Karthikumar S, Jegatheesan K. Isolation of potential antibacterial and antioxidant compounds from *Acalypha indica* and *Ocimum basilicum*. *J Med Plants Res.* 2009; 3(10): 703-706.
- [86] De Sousa MB, Melo OF, De Castro CM, Ximenes EA, Da Silva NH, Da Silva M. Evaluation of the gastric antiulcer, antimicrobial and antioxidant activities of the essential oil from *Ocimum minimum* Linn. *Acta Farm Bonaerense.* 2005; 24(3): 331-336.
- [87] Lukmanul Hakkim F, Arivazhagan G, Boopathy R. Antioxidant property of selected *Ocimum* species and their secondary metabolite content. *J Med Plants Res.* 2008; 2(9): 250-257.
- [88] Dumitriu B, Olariu L, Nita R, Zglimbea L, Rosoiu N. Vascular anti-inflammatory effects of natural compounds from *Aesculus hippocastanum* and *Hedera helix*. *Rom Biotech Lett.* 2013; 18(1): 7963-7974.
- [89] Rai A. The antiinflammatory and antiarthritic properties of ethanol extract of *Hedera helix*. *Indian J Pharm Sci.* 2013; 75(1): 99-102.
- [90] Gülçin İ, Mshvildadze V, Gepdiremen A, Elias R. Antioxidant activity of saponins isolated from ivy:  $\alpha$ -hederin, hederasaponin C, hederacolchiside E and hederacolchiside F. *Planta Med.* 2004; 70(06): 561-563.
- [91] Uddin G, Rauf A, Qaisar M, Latif A, Ali M. Preliminary phytochemical screening and antimicrobial activity of *Hedera helix*. *Middle-East J Sci Res.* 2011; 8(1): 198-202.
- [92] Shi G, Wang B, Wu Q, Wang TC, Wang Cl, Sun XH, Zong WT, Yan M, Zhao QC, Chen YF. Evaluation of the wound-healing activity and anti-inflammatory activity of aqueous extracts from *Acorus calamus* L. *Pak J Pharm Sci.* 2014; 27: 91-95.
- [93] Raja AE, Vijayalakshmi M, Devalarao G. *Acorus calamus* Linn.: chemistry and biology. *Res J Pharm Technol.* 2009; 2(2): 256-261.
- [94] Palani S, Raja S, Praveen R, Devi K, Sivaraj A. Therapeutic efficacy of antihepatotoxic and antioxidant activities of *Acorus calamus* on acetaminophen-induced toxicity in rat. *Int J Integr Biol.* 2009; 7(1): 39-44.
- [95] Fazal H, Ahamad N, Khan MA. Physicochemical, phytochemical evaluation and DPPH-scavenging antioxidant potential in medicinal plants used for herbal

- formulation in Pakistan. *Pak J Bot.* 2011; 43: 63-67.
- [96] Devi SA, Ganjewala D. Antioxidant activities of methanolic extracts of sweet-flag (*Acorus calamus*) leaves and rhizomes. *J Herbs Spices Med Plants.* 2011; 17(1): 1-11.
- [97] Bharathi B, Sundari K, Swamidoss Daniel G. Effect of some medicinal plants on opportunistic bacterial and fungal pathogens associated with HIV. *Asian J Pharm Clin Res.* 2011; 4(1): 149-154.
- [98] Ghosh S, Samanta A, Mandal NB, Bannerjee S, Chattopadhyay D. Evaluation of the wound healing activity of methanol extract of *Pedilanthus tithymaloides* (L.) Poit leaf and its isolated active constituents in topical formulation. *J Ethnopharmacol.* 2012; 142: 714-722.
- [99] Li K, Diao Y, Zhang H, Wang S, Zhang Z, Yu B, Huang S, Yang H. Tannin extracts from immature fruits of *Terminalia chebula* Fructus Retz. promote cutaneous wound healing in rats. *BMC Complement Altern Med.* 2011; 11(86): 1-9.
- [100] Thang PT, Patrick S, Teik LS, Yung CS. Anti-oxidant effects of the extracts from the leaves of *Chromolaena odorata* on human dermal fibroblasts and epidermal keratinocytes against hydrogen peroxide and hypoxanthine-xanthine oxidase induced damage. *Burns.* 2001; 27: 319-327.
- [101] Oberyszyn TM. Inflammation and wound healing. *Front Biosci.* 2007; 12: 2993-99.
- [102] Eming SA, Krieg T, Davidson JM. Inflammation in wound repair: molecular and cellular mechanisms. *J Invest Dermatol.* 2007; 127: 514-525.
- [103] Nonato FR, Santana DG, de Melo FM, dos Santos GGL, Brustolim D, Camargo EA, de Sousa DP, Soares MBP, Villarreal CF. Anti-inflammatory properties of rose oxide. *Int Immunopharmacol.* 2012; 14(4): 779-784.
- [104] Chandran PK, Kuttan R. Effect of *Calendula officinalis* flower extract on acute phase proteins, antioxidant defense mechanism and granuloma formation during thermal burns. *J Clin Biochem Nutr.* 2008; 43: 58-64.
- [105] Kahkeshani N, Farahanikia B, Mahdavian P, Abdolghaffari A, Hassanzadeh G, Abdollahi M, Khanavi M. Antioxidant and burn healing potential of *Galium odoratum* extracts. *Res Pharm Sci.* 2013; 8(3): 197-203.
- [106] Annan K, Houghton PJ. Antibacterial, antioxidant and fibroblast growth stimulation of aqueous extracts of *Ficus asperifolia* Miq. and *Gossypium arboretum* L., wound-healing plants of Ghana. *J Ethnopharmacol.* 2008; 119: 141-144.
- [107] Kumari P, Yadav P, Verma PR, Kumar S, Arya A. A review on wound healing properties of Indian medicinal plants. *Indian J Fundament App Life Sci.* 2013; 3(1): 220-232.
- [108] Nayak BS, Pereira LMP. *Catharanthus roseus* flower extract has wound-healing activity in Sprague Dawley rats. *BMC Complement Altern Med.* 2006; 6(41): 1-6.